

# C2 TRIGONOMETRY

## Worksheet E

- 1 Find all values of  $x$  in the interval  $0 \leq x \leq 360^\circ$  such that
- |   |  |   |                                  |
|---|--|---|----------------------------------|
| <b>a</b> $\sin x = \frac{1}{2}$         | <b>b</b> $\tan x = \sqrt{3}$           | <b>c</b> $\cos x = 0$                   | <b>d</b> $\sin x = -1$           |
| <b>e</b> $\cos x = \frac{\sqrt{3}}{2}$  | <b>f</b> $\sin x = \frac{1}{\sqrt{2}}$ | <b>g</b> $\tan x = -1$                  | <b>h</b> $\cos x = -\frac{1}{2}$ |
| <b>i</b> $\sin x = -\frac{\sqrt{3}}{2}$ | <b>j</b> $\tan x = \frac{1}{\sqrt{3}}$ | <b>k</b> $\cos x = -\frac{1}{\sqrt{2}}$ | <b>l</b> $\tan x = -\sqrt{3}$    |
- 2 Solve each equation for  $\theta$  in the interval  $0 \leq \theta \leq 360^\circ$  giving your answers to 1 decimal place.
- |                                |                                |                                 |                                 |
|--------------------------------|--------------------------------|---------------------------------|---------------------------------|
| <b>a</b> $\cos \theta = 0.4$   | <b>b</b> $\sin \theta = 0.27$  | <b>c</b> $\tan \theta = 1.6$    | <b>d</b> $\sin \theta = 0.813$  |
| <b>e</b> $\tan \theta = 0.1$   | <b>f</b> $\cos \theta = 0.185$ | <b>g</b> $\sin \theta = -0.6$   | <b>h</b> $\tan \theta = -0.7$   |
| <b>i</b> $\cos \theta = -0.39$ | <b>j</b> $\tan \theta = -3.4$  | <b>k</b> $\cos \theta = -0.636$ | <b>l</b> $\sin \theta = -0.203$ |
- 3 Solve each equation for  $x$  in the interval  $0 \leq x \leq 360$ .  
Give your answers to 1 decimal place where appropriate.
- |  |                                       |                                       |
|--|---------------------------------------|---------------------------------------|
| <b>a</b> $\sin(x - 60)^\circ = 0.5$        | <b>b</b> $\tan(x + 30)^\circ = 1$     | <b>c</b> $\cos(x - 45)^\circ = 0.2$   |
| <b>d</b> $\tan(x + 30)^\circ = 0.78$       | <b>e</b> $\cos(x + 45)^\circ = -0.5$  | <b>f</b> $\sin(x - 60)^\circ = -0.89$ |
| <b>g</b> $\cos(x + 45)^\circ = 0.9$        | <b>h</b> $\sin(x + 30)^\circ = 0.14$  | <b>i</b> $\cos(x - 60)^\circ = 0.6$   |
| <b>j</b> $\sin(x - 30)^\circ = -0.3$       | <b>k</b> $\tan(x - 60)^\circ = -1.26$ | <b>l</b> $\sin 2x^\circ = 0.5$        |
| <b>m</b> $\cos 2x^\circ = 0.64$            | <b>n</b> $\sin 2x^\circ = -0.18$      | <b>o</b> $\tan 2x^\circ = -2.74$      |
| <b>p</b> $\sin \frac{1}{2}x^\circ = 0.703$ | <b>q</b> $\tan 3x^\circ = 0.591$      | <b>r</b> $\cos 2x^\circ = -0.415$     |
- 4 Solve each equation for  $x$  in the interval  $0 \leq x \leq 2\pi$  giving your answers in terms of  $\pi$ .
- |   |  |  |
|---|--|--|
| <b>a</b> $\sin x = 0$                                   | <b>b</b> $\cos x = \frac{1}{2}$                  | <b>c</b> $\tan x = 1$                                    |
| <b>d</b> $\cos x = -1$                                  | <b>e</b> $\tan x = -\frac{1}{\sqrt{3}}$          | <b>f</b> $\sin x = -\frac{1}{\sqrt{2}}$                  |
| <b>g</b> $\tan(x + \frac{\pi}{6}) = \sqrt{3}$           | <b>h</b> $\sin(x - \frac{\pi}{4}) = \frac{1}{2}$ | <b>i</b> $\cos(x + \frac{\pi}{3}) = -\frac{\sqrt{3}}{2}$ |
| <b>j</b> $\sin(x + \frac{\pi}{3}) = \frac{1}{\sqrt{2}}$ | <b>k</b> $\cos 2x = -\frac{1}{\sqrt{2}}$         | <b>l</b> $\tan 3x = \frac{1}{\sqrt{3}}$                  |
- 5 Solve each equation for  $\theta$  in the interval  $-180^\circ \leq \theta \leq 180^\circ$ .  
Give your answers to 1 decimal place where appropriate.
- |   |  |   |
|---|--|---|
| <b>a</b> $\cos \theta = 0$                  | <b>b</b> $\tan 2\theta + 1 = 0$              | <b>c</b> $\sin(\theta + 60^\circ) = 0.291$  |
| <b>d</b> $2 \tan(\theta - 15^\circ) = 3.7$  | <b>e</b> $\sin 2\theta - 0.3 = 0$            | <b>f</b> $4 \cos 3\theta = 2$               |
| <b>g</b> $1 + \sin(\theta + 110^\circ) = 0$ | <b>h</b> $5 \cos(\theta - 27^\circ) = 3$     | <b>i</b> $7 - 3 \tan \theta = 0$            |
| <b>j</b> $3 + 8 \cos 2\theta = 0$           | <b>k</b> $2 + 6 \tan(\theta + 92^\circ) = 0$ | <b>l</b> $1 - 4 \sin \frac{1}{3}\theta = 0$ |

## C2 TRIGONOMETRY

## Worksheet E continued

- 6 Solve each equation for  $x$  in the interval  $0 \leq x \leq 180^\circ$ .  
Give your answers to 1 decimal place where appropriate.
- a**  $\tan(2x + 30^\circ) = 1$       **b**  $\sin(2x - 15^\circ) = 0$       **c**  $\cos(2x + 70^\circ) = 0.5$   
**d**  $\sin(2x + 210^\circ) = 0.26$       **e**  $\cos(2x - 38^\circ) = -0.64$       **f**  $\tan(2x - 56^\circ) = -0.32$   
**g**  $\cos(3x - 24^\circ) = 0.733$       **h**  $\tan(3x + 60^\circ) = -1.9$       **i**  $\sin(\frac{1}{2}x + 18^\circ) = 0.572$
- 7 Solve each equation for  $x$  in the interval  $0 \leq x \leq 2\pi$ , giving your answers to 2 decimal places.
- a**  $\tan x = 0.52$       **b**  $\cos 2x = 0.315$       **c**  $\sin(x + \frac{\pi}{4}) = 0.7$   
**d**  $3 \cos x + 1 = 0$       **e**  $\sin \frac{1}{2}x = 0.09$       **f**  $\tan 2x = -0.225$   
**g**  $3 - 4 \sin(x - \frac{\pi}{3}) = 0$       **h**  $\tan(2x + \frac{\pi}{6}) = 2$       **i**  $\cos 3x = -0.81$   
**j**  $5 + 3 \tan x = 0$       **k**  $\cos(2x - \frac{\pi}{2}) = -0.34$       **l**  $1 + 6 \sin 2x = 0$
- 8 **a** Solve the equation  

$$2y^2 - 3y + 1 = 0.$$
**b** Hence, find the values of  $x$  in the interval  $0 \leq x \leq 360^\circ$  for which  

$$2 \sin^2 x - 3 \sin x + 1 = 0.$$
- 9 Solve each equation for  $\theta$  in the interval  $0 \leq \theta \leq 360$ .  
Give your answers to 1 decimal place where appropriate.
- a**  $\sin^2 \theta^\circ = 0.75$       **b**  $1 - \tan^2 \theta^\circ = 0$   
**c**  $2 \cos^2 \theta^\circ + \cos \theta^\circ = 0$       **d**  $\sin \theta^\circ(4 \cos \theta^\circ - 1) = 0$   
**e**  $4 \sin \theta^\circ = \sin \theta^\circ \tan \theta^\circ$       **f**  $(2 \cos \theta^\circ - 1)(\cos \theta^\circ + 1) = 0$   
**g**  $\tan^2 \theta^\circ - 3 \tan \theta^\circ + 2 = 0$       **h**  $3 \sin^2 \theta^\circ - 7 \sin \theta^\circ + 2 = 0$   
**i**  $\tan^2 \theta^\circ - \tan \theta^\circ = 6$       **j**  $6 \cos^2 \theta^\circ - \cos \theta^\circ - 2 = 0$   
**k**  $4 \sin^2 \theta^\circ + 3 = 8 \sin \theta^\circ$       **l**  $\cos^2 \theta^\circ + 2 \cos \theta^\circ - 1 = 0$   
**m**  $\tan^2 \theta^\circ + 3 \tan \theta^\circ - 1 = 0$       **n**  $3 \sin^2 \theta^\circ + \sin \theta^\circ = 1$
- 10 **a** Sketch the curve  $y = \cos x^\circ$  for  $x$  in the interval  $0 \leq x \leq 360$ .  
**b** Sketch on the same diagram the curve  $y = \cos(x + 90)^\circ$  for  $x$  in the interval  $0 \leq x \leq 360$ .  
**c** Using your diagram, find all values of  $x$  in the interval  $0 \leq x \leq 360$  for which  

$$\cos x^\circ = \cos(x + 90)^\circ.$$
- 11 **a** Sketch the curves  $y = \cos x^\circ$  and  $y = \cos 3x^\circ$  on the same set of axes for  $x$  in the interval  $0 \leq x \leq 360$ .  
**b** Solve, for  $x$  in the interval  $0 \leq x \leq 360$ , the equation  

$$\cos x^\circ = \cos 3x^\circ.$$
**c** Hence solve, for  $x$  in the interval  $0 \leq x \leq 180$ , the equation  

$$\cos 2x^\circ = \cos 6x^\circ.$$

# C2 TRIGONOMETRY

## Worksheet F

- 1 a Given that  $4 \sin x + \cos x = 0$ , show that  $\tan x = -\frac{1}{4}$ .
- b Hence, find the values of  $x$  in the interval  $0 \leq x \leq 360^\circ$  for which  $4 \sin x + \cos x = 0$ , giving your answers to 1 decimal place.
- 2 a Show that  $5 \sin^2 x + 5 \sin x + 4 \cos^2 x \equiv \sin^2 x + 5 \sin x + 4$ .
- b Hence, find the values of  $x$  in the interval  $0 \leq x \leq 360^\circ$  for which  $5 \sin^2 x + 5 \sin x + 4 \cos^2 x = 0$ .
- 3 Solve each equation for  $x$  in the interval  $0 \leq x \leq 360^\circ$ . Give your answers to 1 decimal place where appropriate.
- |  |  |
|--|--|
| a $2 \sin x - \cos x = 0$                  | b $3 \sin x = 4 \cos x$                    |
| c $\cos^2 x + 3 \sin x - 3 = 0$            | d $3 \cos^2 x - \sin^2 x = 2$              |
| e $2 \sin^2 x + 3 \cos x = 3$              | f $3 \cos^2 x = 5(1 - \sin x)$             |
| g $3 \sin x \tan x = 8$                    | h $\cos x = 3 \tan x$                      |
| i $3 \sin^2 x - 5 \cos x + 2 \cos^2 x = 0$ | j $2 \sin^2 x + 7 \sin x - 2 \cos^2 x = 0$ |
| k $3 \sin x - 2 \tan x = 0$                | l $\sin^2 x - 9 \cos x - \cos^2 x = 5$     |
- 4 Solve each equation for  $\theta$  in the interval  $-\pi \leq \theta \leq \pi$  giving your answers in terms of  $\pi$ .
- |   |   |
|---|---|
| a $4 \cos^2 \theta = 1$                                   | b $4 \sin^2 \theta + 4 \sin \theta + 1 = 0$           |
| c $\cos^2 \theta + 2 \cos \theta - 3 = 0$                 | d $3 \sin^2 \theta - \cos^2 \theta = 0$               |
| e $4 \sin^2 \theta - 5 \sin \theta + 2 \cos^2 \theta = 0$ | f $\sin^2 \theta - 3 \cos \theta - \cos^2 \theta = 2$ |
- 5 Prove that
- |  |   |
|--|---|
| a $(\sin x + \cos x)^2 \equiv 1 + 2 \sin x \cos x$               | b $\frac{1}{\cos x} - \cos x \equiv \sin x \tan x, \cos x \neq 0$             |
| c $\frac{\cos^2 x}{1 - \sin x} \equiv 1 + \sin x, \sin x \neq 1$ | d $\frac{1 + \sin x}{\cos x} \equiv \frac{\cos x}{1 - \sin x}, \cos x \neq 0$ |
- 6 a Prove the identity  $(\cos x - \tan x)^2 + (\sin x + 1)^2 \equiv 2 + \tan^2 x$ .
- b Hence find, in terms of  $\pi$ , the values of  $x$  in the interval  $0 \leq x \leq 2\pi$  such that  $(\cos x - \tan x)^2 + (\sin x + 1)^2 = 3$ .
- 7  $f(x) \equiv \cos^2 x + 2 \sin x, 0 \leq x \leq 2\pi$ .
- a Prove that  $f(x)$  can be expressed in the form  $f(x) = 2 - (\sin x - 1)^2$ .
- b Hence deduce the maximum value of  $f(x)$  and the value of  $x$  for which this occurs.