

C1 INTEGRATION

Worksheet A

1 Integrate with respect to x

a x^2 **b** x^6 **c** x **d** x^{-4} **e** 5 **f** $3x^2$
g $4x^7$ **h** $6x^{-2}$ **i** $8x^5$ **j** $\frac{1}{3}x$ **k** $2x^{-9}$ **l** $\frac{3}{4}x^{-3}$

2 Find

a $\int (2x + 3) dx$ **b** $\int (12x^3 - 4x) dx$ **c** $\int (7 - x^2) dx$ **d** $\int (x^2 + x + 1) dx$
e $\int (x^4 + 5x^2) dx$ **f** $\int x(x^2 - 3) dx$ **g** $\int (x - 2)^2 dx$ **h** $\int (3x^4 + x^2 - 6) dx$
i $\int (2 + \frac{1}{x^2}) dx$ **j** $\int (x - \frac{1}{x^3}) dx$ **k** $\int x^2(\frac{2}{x^4} - 3) dx$ **l** $\int (x - \frac{4}{x})^2 dx$

3 Integrate with respect to y

a $y^{\frac{1}{2}}$ **b** $y^{\frac{5}{2}}$ **c** $y^{-\frac{1}{2}}$ **d** $4y^{\frac{1}{3}}$ **e** $y^{\frac{3}{4}}$ **f** $5y^{-\frac{2}{3}}$
g $\sqrt[4]{y}$ **h** $\frac{7}{\sqrt{y}}$ **i** $\frac{1}{2y^2}$ **j** $\sqrt{y^3}$ **k** $\frac{5}{2y^4}$ **l** $\frac{1}{3\sqrt{y}}$

4 Find

a $\int (3t^{\frac{1}{2}} - 1) dt$ **b** $\int (2r + \sqrt{r}) dr$ **c** $\int (3p - 1)^2 dp$ **d** $\int (4x + x^{\frac{1}{3}}) dx$
e $\int (\frac{1}{y^3} + y) dy$ **f** $\int (\frac{1}{2}x^2 - x^{\frac{3}{2}}) dx$ **g** $\int \frac{t^3 + 2t}{t} dt$ **h** $\int (r^{\frac{5}{3}} - r^{\frac{2}{3}}) dr$
i $\int \frac{4p^4 - p^2}{2p} dp$ **j** $\int (4 - y^{\frac{7}{4}}) dy$ **k** $\int \frac{1 + 6x^2}{3x^2} dx$ **l** $\int \frac{2t + 3}{\sqrt{t}} dt$

5 Find $\int y dx$ when

a $y = 3x^2 - x + 6$ **b** $y = x^6 - x^3 + 2x - 5$ **c** $y = x(x - 2)(x + 1)$
d $y = (x^{\frac{1}{2}} + 2)^2$ **e** $y = (x^2 - 4)(2x + 3)$ **f** $y = x^3 - 2x^{\frac{4}{3}} + \frac{7}{x^2}$
g $y = \frac{1}{4x^3} - \frac{2}{3x^2}$ **h** $y = (1 - \frac{2}{x^2})^2$ **i** $y = (x^{\frac{5}{2}} - 1)(x^{\frac{3}{2}} + 1)$

6 Find a general expression for y given that

a $\frac{dy}{dx} = 8x + 3$ **b** $\frac{dy}{dx} = \frac{1}{2}x^3 - x^2$ **c** $\frac{dy}{dx} = \frac{4}{3x^3}$
d $\frac{dy}{dx} = (x + 1)^3$ **e** $\frac{dy}{dx} = 2x - \frac{3}{\sqrt{x}}$ **f** $\frac{dy}{dx} = x^{\frac{3}{2}} - 2x^{-\frac{3}{2}}$
g $\frac{dy}{dx} = \frac{3 - x^2}{2x^2}$ **h** $\frac{dy}{dx} = \frac{2}{x^3}(5 - x)$ **i** $\frac{dy}{dx} = \frac{9x - 2}{3\sqrt{x}}$

C1 INTEGRATION

Worksheet B

- 1 a Find $\int (2x + 1) dx$.
- b Given that $\frac{dy}{dx} = 2x + 1$ and that $y = 5$ when $x = 1$, find an expression for y in terms of x .
- 2 Use the given boundary conditions to find an expression for y in each case.
- a $\frac{dy}{dx} = 3 - 6x$, $y = 1$ at $x = 2$ b $\frac{dy}{dx} = 3x^2 - x$, $y = 41$ at $x = 4$
- c $\frac{dy}{dx} = x^2 + 4x + 1$, $y = 4$ at $x = -3$ d $\frac{dy}{dx} = 7 - 5x - x^3$, $y = 0$ at $x = 2$
- e $\frac{dy}{dx} = 8x - \frac{2}{x^2}$, $y = -1$ at $x = \frac{1}{2}$ f $\frac{dy}{dx} = 3 - \sqrt{x}$, $y = 8$ at $x = 4$
- 3 The curve $y = f(x)$ passes through the point $(3, 5)$.
Given that $f'(x) = 3 + 2x - x^2$, find an expression for $f(x)$.
- 4 Given that
- $$\frac{dy}{dx} = 10x^{\frac{3}{2}} - 2x^{-\frac{1}{2}},$$
- and that $y = 7$ when $x = 0$, find the value of y when $x = 4$.
- 5 The curve $y = f(x)$ passes through the point $(-1, 4)$. Given that $f'(x) = 2x^3 - x - 8$,
- a find an expression for $f(x)$,
- b find an equation of the tangent to the curve at the point on the curve with x -coordinate 2.
- 6 The curve $y = f(x)$ passes through the origin.
Given that $f'(x) = 3x^2 - 8x - 5$, find the coordinates of the other points where the curve crosses the x -axis.
- 7 Given that
- $$\frac{dy}{dx} = 3x + \frac{2}{x^2},$$
- a find an expression for y in terms of x .
Given also that $y = 8$ when $x = 2$,
- b find the value of y when $x = \frac{1}{2}$.
- 8 The curve C with equation $y = f(x)$ is such that
- $$\frac{dy}{dx} = 3x^2 + kx,$$
- where k is a constant.
Given that C passes through the points $(1, 6)$ and $(2, 1)$,
- a find the value of k ,
- b find an equation of the curve.

C1 INTEGRATION*Worksheet C*

1 Find

$$\int (x^2 + 6\sqrt{x} - 3) \, dx. \quad (3)$$

2 The curve $y = f(x)$ passes through the point $(1, -2)$.

Given that

$$f'(x) = 1 - \frac{6}{x^3},$$

a find an expression for $f(x)$. (4)The point A on the curve $y = f(x)$ has x -coordinate 2.b Show that the normal to the curve $y = f(x)$ at A has the equation

$$16x + 4y - 19 = 0. \quad (5)$$

3 The curve $y = f(x)$ passes through the point $(3, 22)$.

Given that

$$f'(x) = 3x^2 + 2x - 5,$$

a find an expression for $f(x)$. (4)

Given also that

$$g(x) = (x + 3)(x - 1)^2,$$

b show that $g(x) = f(x) + 2$, (3)c sketch the curves $y = f(x)$ and $y = g(x)$ on the same set of axes. (3)

4 Given that

$$y = x^2 - \frac{3}{x^2},$$

find

a $\frac{dy}{dx}$, (2)b $\int y \, dx$. (3)5 The curve C with equation $y = f(x)$ is such that

$$\frac{dy}{dx} = 3x^2 - 4x - 1.$$

Given that the tangent to the curve at the point P with x -coordinate 2 passes through the origin, find an equation for the curve. (7)6 A curve with equation $y = f(x)$ is such that

$$\frac{dy}{dx} = 3\sqrt{x} - \frac{2}{\sqrt{x}}, \quad x > 0.$$

a Find the gradient of the curve at the point where $x = 2$, giving your answer in its simplest form. (2)Given also that the curve passes through the point $(4, 7)$,b find the y -coordinate of the point on the curve where $x = 3$, giving your answer in the form $a\sqrt{3} + b$, where a and b are integers. (6)

C1 INTEGRATION*Worksheet C continued***7** Find

a $\int (x + 2)^2 dx,$ (3)

b $\int \frac{1}{4\sqrt{x}} dx.$ (3)

8 The curve C has the equation $y = f(x)$ and crosses the x -axis at the point $P(-2, 0)$.

Given that

$$f'(x) = 3x^2 - 2x - 3,$$

a find an expression for $f(x)$, (4)**b** show that the tangent to the curve at the point where $x = 1$ has the equation

$$y = 5 - 2x. \quad (3)$$

9 Given that

$$\frac{dy}{dx} = 2x - \frac{3}{x^2}, \quad x \neq 0,$$

and that $y = 0$ at $x = 1$,**a** find an expression for y in terms of x , (4)**b** show that for all non-zero values of x

$$x^2 \frac{d^2y}{dx^2} - 2y = k,$$

where k is a constant to be found. (4)**10** Integrate with respect to x

a $\frac{1}{x^3},$ (2)

b $\frac{(x-1)^2}{\sqrt{x}}.$ (5)

11 The curve $y = f(x)$ passes through the point $(2, -5)$.

Given that

$$f'(x) = 4x^3 - 8x,$$

a find an expression for $f(x)$, (4)**b** find the coordinates of the points where the curve crosses the x -axis. (4)**12** The curve C with equation $y = f(x)$ is such that

$$\frac{dy}{dx} = k - x^{-\frac{1}{2}}, \quad x > 0,$$

where k is a constant.Given that C passes through the points $(1, -2)$ and $(4, 5)$,**a** find the value of k , (5)**b** show that the normal to C at the point $(1, -2)$ has the equation

$$x + 2y + 3 = 0. \quad (4)$$